

Background and Purpose—We aim to compare the risk of 1-year ischemic stroke recurrence and death for atrial fibrillation diagnosed after stroke (AFDAS), atrial fibrillation known before stroke (KAF), and sinus rhythm (SR).

Methods—From June 2012 to January 2013, 6004 patients with acute ischemic stroke were admitted to 219 urban hospitals in the China National Stroke Registry II. Based on heart rhythm assessed during admission, we classified patients as AFDAS, KAF, or SR. We explored the relationship between heart rhythm groups and 1-year ischemic stroke recurrence or death by using Cox regression adjusted for multiple covariates. Considering that death is a competing risk for stroke recurrence, we used the competing risks analysis of Fine and Gray and subdistribution Cox proportional hazards to test the association between heart rhythm and 1-year outcomes.

Results—Among 19604 ischemic stroke patients, 7127 had SR, 495 AFDAS, and 1382 KAF. At 1 year, 54 (10.9%) patients with AFDAS, 182 (13.2%) with KAF, and 1008 (5.7%) with SR had recurrent ischemic strokes ($P < 0.0001$). Mortality was 22.0% in patients with AFDAS, 22.1% in patients with KAF, and 7.0% in patients with SR ($P < 0.0001$). AFDAS-related ischemic stroke recurrence adjusted risk was higher than that of SR (adjusted subdistribution hazard ratio 1.61; 95% CI 1.17–2.21).



Stroke

A

with minimal or no participation of central neurogenic mechanisms, AFDAS may comprise a heterogeneous mix of AF phenotypes, possibly including cardiogenic and neurogenic types, as well as a combination of both. Several studies have explored the underlying pathophysiologic mechanisms of AFDAS by comparing the cardiovascular risk factors as well as cardiac structure and function between KAF and AFDAS, though the results were not consistent.^{6,7} According to a previous study, stroke patients with KAF and AFDAS share common cardiovascular risk factors,

⁴ Compared

with KAF, which is presumed to be primarily cardiogenic

similar echocardiographic findings, and suffer equally severe strokes.⁶ Another study showed that patients with AFDAS have a low frequency of underlying cardiac disease and a strikingly high proportion of concurrent strategic insular infarctions, which suggests that neurogenic mechanisms are responsible. Recently, a study found that the AFDAS and sinus rhythm (SR) have similar 1-year ischemic stroke recurrence, suggesting that the underlying pathophysiology of AFDAS may differ from that of KAF.⁷

Considering the inconsistency of the previous studies, we aim to further characterize the clinical and prognostic features of AFDAS and compare them with those of KAF and SR in our study. Patients with AFDAS will comprise a mix of AFDAS phenotypes: neurogenic with potentially better outcomes and cardiogenic with worse outcomes. We aim to compare the prevalence of cardiovascular comorbidities and prognosis between AFDAS, KAF, and SR to further explore the pathophysiology of AFDAS.

Methods

The data that support the findings of this study are available from the corresponding author on reasonable request.

Study Cohort

China National Stroke Registry II (CNSR II), launched in 2012 by the Ministry of Health of China, is a nationwide initiative to establish a reliable national stroke database for evaluating the delivery of stroke care in clinical practice. The criteria for site selection in China National Stroke Registry I (CNSR I) have been published in order for the hospital characteristics to be similar between CNSR I and CNSR II, the same criteria were used for selection. Patients were recruited consecutively from all 219 urban hospitals voluntarily participating in the China National Stroke Registry II from June 2012 to January 2013 if they met the following criteria: (1) age > 18 years; (2) diagnosis within 7 days of the index event of ischemic stroke, TIA, spontaneous intracerebral hemorrhage, or subarachnoid hemorrhage; (3) direct hospital admission from a physician's clinic or emergency department; and (4) informed consent provided by the patient or a legally authorized representative.

The study was approved by the central Institutional Review Board at Beijing Tiantan Hospital. Every participant provided written informed consent before participation.

Data Collection and Management

Patient information including demographics, risk factors, comorbidities, medications, selected laboratory tests, discharge status, and hospital-level characteristics were systematically collected during hospitalization and at discharge by trained research coordinators at each participating hospital. The severity of stroke was assessed by the National Institutes of Health Stroke Scale. Stroke risk was calculated at discharge using the CHA₂DS₂-VASc score (congestive heart failure, hypertension, age 65–74 years and age 75 years, diabetes mellitus, prior stroke or TIA; vascular disease, and female sex). Based on this definition, all patients had minimum CHA₂DS₂-VASc risk scores of 2 due to their index stroke.

Eligibility and Criteria

Patients underwent the 12-lead ECG or further 24-hour Holter cardiac rhythm recording/telemetry at the time of admission or during the hospitalization based on the discretion of the treating physician. Based on their history of AF and on the results of ECG monitoring during hospitalization, patients were stratified into 3 main groups according to a previous study¹ (1) AFDAS (no history of AF but AF detected during hospital stay after the index ischemic stroke), (2)

KAF (history of AF known before the index ischemic stroke, regardless of whether it was documented during admission), and (3) SR (no history of AF and no AF identified during admission).

Outcome Measures

The primary outcome of this study was the risk of a recurrent ischemic stroke and death at 1 year after discharge, which was ascertained by using a previously validated algorithm. Patients were contacted by telephone at 3, 6, and 12 months after stroke onset by trained research personnel who followed standard scripts to collect information on death, stroke recurrence, and disability. During the follow-up, stroke recurrences associated with rehospitalization were sourced to the corresponding hospitals to ensure reliable diagnosis. In case of a suspected recurrent cerebrovascular event without hospitalization, adjudication was performed by the research coordinators together with the principal investigator. The detailed follow-up procedure has been described previously.¹

Statistical Analysis

Continuous variables are reported as mean with SD or median with interquartile ranges, and categorical variables are reported as frequencies with percentages. We adopted cumulative incidence function curves for recurrent ischemic stroke in each of the 3 heart rhythm groups, and comparison was conducted using the Fine and Gray test. We estimated the association between each AF subgroup and recurrent ischemic stroke at 1 year after discharge by using subdistribution Cox proportional hazards. We developed 2 different models by using variables known to influence the risk of recurrent ischemic stroke. Model 1 was adjusted for age groups, sex, stroke severity, and prescription of anticoagulants at discharge. Model 2 was adjusted for age groups, sex, stroke severity, systemic hypertension, diabetes mellitus, congestive heart failure, coronary artery disease, peripheral artery disease, myocardial infarction, prior history of stroke or TIA, and prescription of anticoagulants at discharge. We used the χ^2 with 2-tailed probabilities for comparing proportions. All tests were 2-tailed with $P < 0.05$ considered statistically significant. All analyses were performed using SAS version 9.4 software (SAS Institute, Cary, NC).

Results

Among 19604 ischemic stroke patients, 495 (2.5%) had AFDAS, 1382 (7.0%) had KAF, and 1727 (9.4%) had SR (Table 1). At 1 year, 1244 (6.3%) patients had a recurrent ischemic stroke: 54 (10.9%) patients with AFDAS, 182 (13.2%) with KAF, and 1008 (5.7%) with SR had recurrent ischemic strokes ($P < 0.0001$).

AFDAS Versus KAF and SR

When comparing characteristics between patients with KAF and AFDAS, fewer patients with AFDAS were female (41.1% versus 48.6%; $P = 0.006$), had a previous stroke or TIA (32.5% versus 45.6%; $P < 0.0001$), or were diagnosed with coronary heart disease (22.8% versus 31.8%; $P = 0.0002$), heart failure, or myocardial infarction (6.9% versus 15.9%; $P < 0.0001$). Compared with patients with SR, those with AFDAS had a significantly higher prevalence of coronary artery disease (22.8% versus 9.8%; $P < 0.0001$), heart failure, or myocardial infarction (6.9% versus 3.3%; $P < 0.0001$; Table 1 in the [online-only Data Supplement](#)).

Risk of Recurrent Ischemic Stroke at 1 Year

Accounting for the competing risk of death, the cumulative incidence of recurrent ischemic stroke within 1 year after discharge for patients with AFDAS was higher than that for participants with SR (Figure 1).



Stroke

Patients with KAF and patients with AFDAS both had a higher risk of death at 1 year (hazard ratio, 1.55; 95% CI, 1.32–1.82; and hazard ratio, 1.70; 95% CI, 1.37–2.12, respectively) than those with SR in all the cause-specific Cox regression models (Table 2). The risk of death at 1 year in patients with AFDAS was not different from that of patients with KAF (hazard ratio, 1.10; 95% CI, 0.86–1.41; Table 2).

Discussion

In our study of 1222 ischemic stroke patients without a history of AF, AFDAS was identified in 495 patients. There are several potential explanations for the lower rate of AF among the ischemic stroke patients in our study. First, the stroke population in our study was younger than that in similar studies in Western countries,^{15,16} and AF-associated stroke is associated with higher age. Furthermore, studies using prolonged noninvasive and invasive cardiac monitoring devices have shown increased detection of AF after ischemic stroke or TIA. In our study, however, AF was diagnosed at discharge according to the 12-lead ECG/24-hour Holter cardiac rhythm recording at the time of admission or during the hospitalization. As such, the detection rate of AF may have been underestimated.

We found a lower frequency of underlying cardiac disease, previous stroke, or TIA and lower CHA2DS2-VASc scores in patients with AFDAS compared with those with KAF, which were consistent with findings by Sposato et al. These results support the hypothesis that AFDAS may comprise a wide spectrum of AF with cardiovascular risk factors that are dissimilar to those of KAF. However, this result was different from findings by Bisson et al,¹⁹ which found that preexisting cardiovascular comorbidities underlie newly

diagnosed poststroke AF. The discrepancy may be attributable to different definitions of AFDAS, KAF, and new AF in these 2 studies. In our study, patients with AFDAS were defined as those with no history of AF and had AF detected during their hospital stay after the index ischemic stroke. In the study by Bisson et al, a medical history of AF or AF recorded during the first 30 days of hospital stay was considered as KAF at baseline. Patients with AFDAS in our study were classified as having KAF in the study by Bisson et al. Furthermore, the definition of patients with new AF in the study by Bisson et al were those without AF at baseline and diagnosed as having incident AF during a subsequent hospitalization after a follow-up period of 7.9±11.5 months. In contrast, we did not have follow-up data for AF.

Considering the low rate of anticoagulant use in patients with AFDAS both at discharge and at 1-year follow-up, the rate of stroke recurrence in AFDAS patients may reflect the natural course of AF that was triggered by the stroke. At 1 year, patients with AFDAS had a higher risk of recurrent ischemic stroke and death than those with SR, even after adjusting for main confounders. When compared with KAF, patients with AFDAS had no difference in the risk of recurrent ischemic stroke or death at 1 year.

Our finding is in line with the study by Lip et al,²¹ which found that among patients with an index stroke or TIA, patients with early AF (AF first diagnosed within 12 months before or within 7 days after the index stroke) had a higher hazard ratio for stroke recurrence within 1 year relative to patients without AF. They also observed that patients without oral anticoagulation prescribed at baseline and having AF first diagnosed >7 days poststroke (late AF) was highly associated with recurrent stroke/TIA. Although our study defined AFDAS as both AF detected within 7 days after index stroke and partial AF diagnosed between 7 days poststroke and discharge from the hospital, both of them were found to have higher stroke recurrence than patients without AF.

Other studies that explored the risk factors for stroke recurrence and mortality after ischemic stroke in patients with AF have found that age, National Institutes of Health Stroke Scale at admission, gender, and the absence of oral anticoagulation were associated with poor outcome or a higher rate of stroke recurrence and mortality.²² In our study, we compared the baseline characteristics between patients with KAF and AFDAS, and we found that there were no differences in the risk factors mentioned above between the 2 groups.

However, our results were not consistent with a recent study by Sposato et al,²⁰ which found that compared with patients with SR, those with AFDAS had no difference in the risk of recurrent ischemic stroke at 1 year. They explained these results by hypothesizing that the presence of a given proportion of relatively benign neurogenic AFDAS phenotypes may dilute the overall AFDAS-associated risk of recurrent ischemic stroke. Indeed, several studies have found possible pathophysiologic mechanisms underlying neurogenic AFDAS such as the involvement of the insular cortex,^{23–25} systemic inflammation,^{26–28} and autonomic dysfunction.²⁹ The finding that patients with KAF have a higher prevalence of cardiovascular risk factors in our study was also compatible with the result by Sposato et al,²⁰ however,

whether neurogenically triggered AFDAS is associated with a lower risk of recurrent stroke than cardiogenic AFDAS caused by underlying heart disease remains uncertain. There are several limitations in our study. First, some patients in the SR group may have had covert paroxysmal AF due to a lack of systematic evaluation of AFDAS, and We were unable to compare the proportion of neurogenic AFDAS between these 2 studies as neither of them can provide adequate distinction between the pathophysiologic mechanisms underlying AFDAS.

Stroke

Conclusions

Compared with SR, AFDAS has a higher risk of 1-year ischemic stroke recurrence and mortality but shares a similar level of risk with KAF. The potential risk of AFDAS should not be overlooked, and it is important to understand the underlying pathophysiology to help improve treatment in the future.

Acknowledgments

We thank all participating hospitals, their physicians, and nurses, and the China National Stroke Registry II Steering Committee members.

Sources of Funding

This work was supported by grants from National Key R&D Program of China (2017YFC1310901, 2016YFC0901001, 2016YFC0901002), Beijing Municipal Committee of Science and Technology (D151100002015003), Beijing Municipal Administration of Hospitals' Mission Plan (SML20150502), the Ministry of Science and Technology of the People's Republic of China (2011BAI08B02, 2012ZX09303, and 2013BAI09B03), Beijing Institute for Brain Disorders (BIBD-PXM2013_014226_07_000084).

Disclosures

None.

References

1. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke*. 1991;22:983–988.
2. Sposato LA, Cipriano LE, Saposnik G, Ruiz Vargas E, Riccio PM, Hachinski V. Diagnosis of atrial fibrillation after stroke and transient ischaemic attack: a systematic review and meta-analysis. *Lancet Neurol*. 2015;14:377–387. doi: 10.1016/S1474-4422(15)70027-X
3. Spence JD. Cardioembolic stroke: everything has changed. *Stroke Vasc Neurol*. 2018;3:76–83. doi: 10.1136/svn-2018-000143
4. Cerasuolo JO, Cipriano LE, Sposato LA. The complexity of atrial fibrillation newly diagnosed after ischemic stroke and transient ischemic attack: advances and uncertainties. *Curr Opin Neurol*. 2017;30:28–37. doi: 10.1097/WCO.0000000000000410
5. Sposato LA, Riccio PM, Hachinski V. Poststroke atrial fibrillation: cause or consequence? Critical review of current video neurology. 2014;82:1180–1186. doi: 10.1212/WNL.0000000000000265
6. Rizos T, Horstmann S, Dittgen F, Täger T, Jenetzky E, Heuschmann P, et al. Preexisting heart disease underlies newly diagnosed atrial fibrillation after acute ischemic stroke. *Stroke*. 2016;47:336–341. doi: 10.1161/STROKEAHA.115.011465
7. González Toledo ME, Klein FR, Riccio PM, Cassará FP, Muñoz Giacomelli F, Racosta JM, et al. Atrial fibrillation detected after acute ischemic stroke: evidence supporting the neurogenic hypothesis. *J Stroke Cerebrovasc Dis*. 2013;22:e486–e491. doi: 10.1016/j.jstrokecerebrovasdis.2013.05.015
8. Sposato LA, Cerasuolo JO, Cipriano LE, Fang J, Fridman S, Paquet M, et al; PARADISE Study Group. Atrial fibrillation detected after stroke is related to a low risk of ischemic stroke recurrence. *Neurology*. 2018;90:e924–e931. doi: 10.1212/WNL.00000000000005126
9. Li Z, Wang C, Zhao X, Liu L, Wang C, Li H, et al; China National Stroke Registries. Substantial progress yet significant opportunity for improvement in stroke care in China. *Stroke*. 2016;47:2843–2849. doi: 10.1161/STROKEAHA.116.014143
10. Wang Y, Cui L, Ji X, Dong Q, Zeng J, Wang Y, et al; China National Stroke Registry Investigators. The China National Stroke Registry for patients with acute cerebrovascular events: design, rationale, and baseline patient characteristics. *Int J Stroke*. 2011;6:355–361. doi: 10.1111/j.1747-4949.2011.00584.x
11. Lyden P, Brott T, Tilley B, Welch KM, Mascha EJ, Levine S, et al. Improved reliability of the NIH Stroke Scale using video training. NINDS TPA Stroke Study Group. *Stroke*. 1994;25:2220–2226.
12. *veiyon brillatiop. Cthet.*
199;94:4963•096.
51. Le EMFe,Grau-Sepul(v)15

